

# **STATISTICS**

Finds and describes patterns in data

"Rob Bob," world's tallest high school basketball player


Algebra 1A \* Ms Johnson

**Austin Horner 1-2023** 











Later, he joined the basketball program at the University of Rochester.

ube.com/watch?v=3\_EuHpeZ9 Qc

https://www.youtube.com/watch?v FUijFFe566c









# **STATISTICS**

Finds and describes patterns in data





## 5 number summary


Rob Bob's 3rd grade class




















How do you make a 5 number summary?




shutterstock.com · 1722790780

First, arrange all the data in ascending order.




shutterstock.com · 1722790780

First, arrange all the data in ascending order.




shutterstock.com · 1722790780


Identify 5 number summary: Minimum and maximum. Median, then Q1 and Q3.



The IQR, the **Interquartile** range is Q3 - Q1. Use all of this info to make a box and whisker plot.



Chartio.com







Probability of rolling 6 twice: (%)(%) = 1/36. Rolling 3 or a 6 is (%) + (%) = 2/6 = 1/3

A statistician is compiling a table summarizing results from a survey about whether seventh and eighth grade students walk or take a bus to school. The statistician gets interrupted before completing the last cell of the table.

	$7^{ m th}~{ m grade}$	$8^{ m th}$ grade
walks	82	96
rides bus	144	

What value should be entered into the blank cell if it is known that 40% of the surveyed eighth grade students ride the bus to school?

Let x be the missing value in the table. Then there are 96+x eighth grade students who were surveyed, of which x of them ride the bus to school. So, the fraction of eighth grade students who ride the bus to

of which 
$$x$$
 of them ride the bus to school. So, the fraction of eighth grade students who ride the bus to school is  $\frac{x}{96+x}$ . This is known to equal  $40\%=0.4$ . Consequently, 
$$\frac{x}{96+x}=0.4$$
 
$$x=0.4\,(96+x)$$


$$x=0.4\left(96\right)+0.4x$$

$$m = 0.1(00) + 0.10$$

$$x - 0.4x = 0.4 (96)$$
$$x - 0.4x = 38.4$$

$$0.6x = 38.4$$
 $x = \frac{38.4}{3.3}$ 

x = 64.










The coefficient that best fits the first graph is -0.65 because the data points are somewhat close together and fit a linear trend with a negative slope.

#### Correlation and Scatter Plots

**Positive Correlation**: as the independent variable increases, the dependent variable also increases.

**Negative Correlation**: as the independent variable decreases, the dependent variable also decreases.

No correlation: there is no correlation, relationship, between the two variables.




### WHAT IS A SCATTER PLOT?




#### **Scatter Plot**

A **scatter plot** is a graph that shows the relationship between two data sets. The two sets of data are graphed as ordered pairs in a

coordinate plane.





Age and Pets Owned

10 20 30 40 50 60 70 x

Person's age (years)



Notes from Ms Johnson class

### Practice Problems - do these along with me!

Remember, the independent and dependent variables. Which one depends on which?

What kind of relationship do these variables typically have?

- Height and shoe size
- Distance driving and amount of gas in tank
- Hours worked and pay
- Video games purchased and your bank account

Notes from Ms Johnson class





# **CREDITS**

StrictlyBBall (2022)
What happened to Robert Bobroczky. <u>Youtube.com</u>

Geogebra.org (2022)
Box plot tool. <a href="https://www.geogebra.org/m/r8axwgzr">https://www.geogebra.org/m/r8axwgzr</a>

Horner, Celeste (2022) Learning coach